

RESEARCH ARTICLE

Prebiotic Potential of a Maize-Based Soluble Fibre and Impact of Dose on the Human Gut Microbiota

Adele Costabile^{1,2*}, Eddie R. Deaville¹, Agustin Martin Morales¹, Glenn R. Gibson¹

1 Department of Food and Nutritional Sciences, University of Reading, Whiteknights, PO Box 226, Reading, RG6 6AP, United Kingdom, **2** Health Sciences Research Centre, Life Sciences Department, Whitelands College, University of Roehampton, London, Holybourne Ave, London, SW15 4JD, United Kingdom

* a.costabile@reading.ac.uk; adele.costabile@roehampton.ac.uk



CrossMark
click for updates

OPEN ACCESS

Citation: Costabile A, Deaville ER, Morales AM, Gibson GR (2016) Prebiotic Potential of a Maize-Based Soluble Fibre and Impact of Dose on the Human Gut Microbiota. PLoS ONE 11(1): e0144457. doi:10.1371/journal.pone.0144457

Editor: Christian U. Riedel, University of Ulm, GERMANY

Received: June 22, 2015

Accepted: November 18, 2015

Published: January 5, 2016

Copyright: © 2016 Costabile et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: The present study was funded by Tate & Lyle Ingredients Americas LLC. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The present study was funded by Tate & Lyle Ingredients Americas LLC. The research and all publications arising out of or referable to it are considered proprietary data to which Tate & Lyle Ingredients Americas LLC claims exclusive right of reference in accordance with

Abstract

Dietary management of the human gut microbiota towards a more beneficial composition is one approach that may improve host health. To date, a large number of human intervention studies have demonstrated that dietary consumption of certain food products can result in significant changes in the composition of the gut microbiota i.e. the prebiotic concept. Thus the prebiotic effect is now established as a dietary approach to increase beneficial gut bacteria and it has been associated with modulation of health biomarkers and modulation of the immune system. Promitor™ Soluble Corn Fibre (SCF) is a well-known maize-derived source of dietary fibre with potential selective fermentation properties. Our aim was to determine the optimum prebiotic dose of tolerance, desired changes to microbiota and fermentation of SCF in healthy adult subjects. A double-blind, randomised, parallel study was completed where volunteers (n = 8/treatment group) consumed 8, 14 or 21 g from SCF (6, 12 and 18 g/fibre delivered respectively) over 14-d. Over the range of doses studied, SCF was well tolerated Numbers of bifidobacteria were significantly higher for the 6 g/fibre/day compared to 12g and 18g/fibre delivered/day (mean 9.25 and 9.73 Log₁₀ cells/g fresh faeces in the pre-treatment and treatment periods respectively). Such a numerical change of 0.5 Log₁₀ bifidobacteria/g fresh faeces is consistent with those changes observed for inulin-type fructans, which are recognised prebiotics. A possible prebiotic effect of SCF was therefore demonstrated by its stimulation of bifidobacteria numbers in the overall gut microbiota during a short-term intervention.

Introduction

The colon contains the most abundant and diverse assemblage of bacteria in the human body [1]. Symbiotic interactions within this complex community are now recognized as important mediators of health by the scientific community [2]. Like other environmental microbial communities, the human microbiome is a complex and dynamic system that plays an important role in many aspects of host physiology [3–6]. Together with the gut immune system, colonic and mucosal microbiota contributes to a barrier that prevents pathogenic bacteria from

Regulation (EC) no. 1924/2006 of the European Parliament and of the Council on Nutrition and Health Claims Made on Foods. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials.

invading the gastrointestinal (GI) tract. Generally, bacteria that have an almost exclusive saccharolytic metabolism can be generally considered as beneficial, e.g. bifidobacteria [7–11]. The intestinal microbiota salvages energy through fermentation of carbohydrates and other substrates not digested in the upper gut [12–14]. In addition, some dietary protein may reach the colon, while endogenous secretions, including mucin, provide diet-independent substrates. Many secondary plant metabolites ingested with the diet, such as polyphenolic substances, may also reach the large intestine and are subject to bacterial transformation. [14]. Colonic bacteria use a range of carbohydrate hydrolyzing enzymes to produce gases and organic acids. Dietary components that stimulate fermentation lead to an increase in bacterial mass and consequently faecal mass and, thus have a stool bulking effect [15]. It is estimated that about 30 g of bacteria are produced for every 100 g of carbohydrate that is fermented. According to Gibson and Roberfroid [16] predominant bacterial groups of the human gut microbiota may be classified into those that are either beneficial with health promoting effects; harmful or pathogenic; or those that demonstrate both or no effects.

Members of nine bacterial phyla were found to inhabit the human gastrointestinal tract, of which *Firmicutes* and *Bacteroidetes* are the dominant ones, while Actinobacteria and Proteobacteria subdominants [4]. Besides the low level of biodiversity at the phylum level, the human intestinal microbiota is tremendously diverse at species and subspecies level [3]. It is characterized by a considerable interpersonal variation in bacterial species and strains [7] and, up to now, 1000 different bacterial species-level phylogenetic types have been identified in the human intestinal microbiota. However, because of a significant degree of functional redundancy, the differences in the microorganism lineage between individuals are greater than differences in the representation of the gene networks embedded in the individual microbiomes. To date, it seems clear that the relative composition of microbes within the gut microbiota is largely reliant upon food consumed, one opportunity for modulate the microbiome is through dietary change [17–20]. For this reason, dietary management of the human gut microbiota towards a more beneficial composition is an approach that can improve host health [21–22]. Bacteria such as bifidobacteria and lactobacilli have demonstrable beneficial effects through improving resistance to gut infections by inhibiting the growth of harmful bacteria [23], protecting against bacterial invasion [24], reducing cholesterol levels [25–26], improving the immune response [27] and producing vitamins within the gut [28].

The scientific understanding of dietary modulation of the human gut microbiota has been developed over several decades, with probiotics as a principal focus [18, 22]. While the role of probiotics has been widely advocated, it is possible that a greater number of food products will further exploit the prebiotic approach over the next few years. Prebiotics can also be added to many foods including those which are cooked or baked, which are more problematic for probiotic enrichment [29–30].

A maize-based soluble fibre, Soluble Corn Fibre (SCF), has been shown to be a versatile source of dietary fibre with potential selective fermentation properties as determined in vitro [31] and in vivo [32]. SCF is a glucose polymer obtained from a partially hydrolyzed starch-made glucose syrup. Possessing a high digestive tolerance after ingestion [33–35], SCF has also been shown to elicit a low post-prandial blood glucose and insulin response (i.e. reduced glycaemic potential) [33]. Recently, SCF has been shown to improve bone properties [36], ameliorate the immune response in experimental colitis [35], and improve laxation [37–38], which is one of the most well noted benefits of dietary fibres. Key to the establishment of novel prebiotic products is assessment of an optimum dose level required to elicit changes in the gut microbiota during feeding. Therefore, the aim of the present pilot human study was to determine the effect of different doses of SCF for prebiotic potential in healthy volunteers and their impact on the human intestinal microbial ecosystem. Stool samples were collected before and after

treatment and changes within the gut microbiota profile were determined. Nine 16S rRNA-based fluorescence in situ hybridization probes were used to target predominant groups/species of human faecal microbiota. Gastrointestinal symptoms and stool characteristics such as stool frequency, consistency, abdominal pain, intestinal bloating and flatulence were also recorded in order to assess prebiotic tolerance.

Materials and Methods

Selection and characteristics of study population

Twenty-four healthy volunteers (gender: 12 female, 12 male; age: mean 33, range 18 to 50 yr; body mass index (BMI): mean 23.6 kg/m², range 20.1 to 29.7 kg/m²) were recruited from the local community. The study was approved by the University of Reading Research Ethics Committee prior to the start of the study and written informed consent was obtained from all subjects prior to commencement of selection. Inclusion criteria were males and females from 18 to 50 years of age and BMI 18.5 to <30 kg/m². General health status was assessed using a pre-study medical questionnaire and putative volunteers were excluded if they fulfilled any of the following exclusion criterion: diagnosed as suffering from any chronic gastrointestinal complaints (including chronic constipation, diarrhoea or irritable bowel syndrome), diabetes or anaemia; requirement to take long-term medication active on the gastro-intestinal tract, for the treatment of cardio-vascular disease, or any other long-term medication; high blood cholesterol or use of cholesterol lowering drugs/functional foods; history of drug or alcohol misuse or alcohol consumption exceeding 14 and 21 units/week for females and males respectively; those suffering with any allergies to medication or food; smokers, and those on weight reducing diets. Females were excluded who were either planning pregnancy within six months of the start of the study, lactating, or had given birth within the preceding six months. Specified timed exclusion criteria included use of antibiotics with the previous six months, participation in any probiotic, prebiotic or laxative study in the previous three months, or intake of an experimental drug within four weeks of the study start. Consumption of any probiotic or prebiotic products and use of medication active on the gastrointestinal tract or use for the treatment of abdominal pain, intestinal discomfort, or constipation were prohibited during the four weeks prior to commencement of the study and for the duration of the study.

Study design and treatments

The overall study design is outlined in [Fig 1](#) and was completed in a double-blind manner, with two experimental periods (pre-treatment and treatment period) each lasting 14 days. Eight volunteers were randomly allocated on the basis of age and BMI to each of three treatment groups. The treatments were designed to provide a total of 8, 14 or 21 g from SCF (Promitor™ Soluble Corn Fibre, SCF) (equivalent to 6, 12 and 18 g/fibre delivered, respectively). Supplements were provided by Tate & Lyle, France, as dry powders and were consumed as 250 ml beverages twice daily. Volunteers were requested to substitute 500 ml of their normal daily fluid intake to compensate for the volume of water consumed with the treatments. Freshly voided faecal samples were provided by each volunteer at the end of the pre-treatment period (baseline sample) and then after 14 days of dietary intervention during the treatment period.

Faecal samples were collected in sterile plastic containers, stored in an anaerobic cabinet (10% H₂; 10% CO₂; 80% N₂) and then processed within two hours of collection.

Throughout the pre-treatment and treatment periods, volunteers recorded details of bowel habits including stool frequency and consistency (Bristol stool scale) [39], stomach or intestinal bloating, abdominal pain, incidence and frequency of flatulence and stomach noises. This




Fig 1. Study design of a double-blind, randomised, parallel study with eight volunteers per treatment group, and consisting of a pre-treatment and treatment period (14-days). Treatments with were 8, 14 and 21 g-SCF/d designed to 6, 12 and 18 g fibre/delivered/day from Promitor™ Soluble Corn Fibre (SCF).

doi:10.1371/journal.pone.0144457.g001

information was recorded as being indicative of digestive tolerance. In addition, volunteers recorded details of dietary intake on four days within each period (data not reported).

Faecal preparation and enumeration of faecal microbial populations by fluorescence *in situ* hybridisation

Freshly voided faecal samples (~ 2–5 g) were diluted 1 in 10 (w/w) with sterile, anaerobic phosphate buffered saline (PBS, 0.1M; pH 7.0) and homogenised using a Stomacher 400® (Seward, Norfolk, UK) for 2 min at normal speed.

Fluorescence *in situ* hybridisation (FISH) was performed as described by Costabile et al. [40]. The composition of hybridisation and wash buffers depends on the rRNA-targeted oligonucleotide probe as reported in probeBase (<http://www.microbial-ecology.net/probebase>) and was used accordingly. All oligonucleotide probes used in the study are reported in Table 1 [40]. Briefly, aliquots (375 µl) of 1 in 10 w/w stool sample suspension in PBS, 0.1M were fixed overnight at 4°C with 4% (w/v) filtered paraformaldehyde (pH 7.2) in a ratio of 1:4 (v/v), washed twice with filtered PBS (0.2 µm pore size), resuspended in 300 ml of a PBS/ethanol mixture (1:1, v/v) and then stored at -20°C for up to 3 months. For the hybridizations, 20 µl of each sample was pipetted onto Teflon and poly-L-lysine-coated, six-well (10 mm diameter each) slides (Tekdon Inc., Myakka City, FL). To permeabilize the cells for use with probe Lab158, samples were treated with 50 µl of lysozyme (1 mg/ml in 100 mM Tris-HCl, pH 8.0) at 37°C for 15 min before being washed (2–3 s) in water and were finally dehydrated in the ethanol series (50%, 80% and 96% v/v ethanol, 3 min each). A probe/hybridization buffer mixture (5 ng/5 µl of a 50 ng/µl probe in stock solution plus 45 µl of hybridization buffer) was applied

Table 1. Probes used for Fluorescence *in situ* hybridization (FISH) analysis of bacterial populations in human faeces.

Short name	Accession no.*	Full name	Target species	Sequence (5' to 3')
Bif164	pB-00037	S-G-Bif-0164-a-A-18	Most <i>Bifidobacterium</i> spp. and <i>Parascardovia denticolens</i>	CATCCGGCATTACCACCC
Lab158	ND	S-G-Lab-0158-a-A-20	Most <i>Lactobacillus</i> , <i>Leuconostoc</i> and <i>Weissella</i> spp.; <i>Lactococcus lactis</i> ; all <i>Vagococcus</i> , <i>Enterococcus</i> , <i>Melissococcus</i> , <i>Tetragenococcus</i> , <i>Catelicoccus</i> , <i>Pediococcus</i> and <i>Paralactobacillus</i> spp.	GGTATTAGCAYCTGTTTCCA
Bac303	pB-00031	S-*-Bacto-0303-a-A-17	Most <i>Bacteroides sensu stricto</i> and <i>Prevotella</i> spp.; all <i>Parabacteroides</i> ; <i>Barnesiella viscericola</i> and <i>Odoribacter splanchnicus</i>	CCAATGTGGGGGACCTT
Chis150	pB-00962	S-*-Chis-0150-a-A-23	Most members of <i>Clostridium</i> cluster I; all members of <i>Clostridium</i> cluster II; <i>Clostridium tyrobutyricum</i> ; <i>Adhaeribacter aquaticus</i> and <i>Flexibacter canadensis</i> (family <i>Flexibacteriaceae</i>); [<i>Eubacterium</i>] <i>combesii</i> (family <i>Propionibacteriaceae</i>)	TTATGCGGTATTAATCTYCCTT
Ato291	pB-00943	S-*-Ato-0291-a-A-17	<i>Atopobium</i> , <i>Colinsella</i> , <i>Olsenella</i> and <i>Eggerthella</i> spp.; <i>Cryptobacterium curtum</i> ; <i>Mycoplasma equigenitalium</i> and <i>Mycoplasma elephantis</i>	GGTCGGTCTCTCAACCC
Erec482	pB-00963	S-*-Erec-0482-a-A-19	Most members of <i>Clostridium</i> cluster XIVa; <i>Syntrophococcus sucromutans</i> , [<i>Bacteroides</i>] <i>galacturonicus</i> and [<i>Bacteroides</i>] <i>xylanolyticus</i> , <i>Lachnospira pectinschiza</i> and <i>Clostridium saccharolyticum</i>	GCTTCTTAGTCARGTACCG
Eub I-II-III	pB-00159-160-161	S-D-Bact-0338-a-A-18	Most domain bacteria	GCT GCC TCC CGT AGG AGT
Rrec584	ND	ND	<i>Roseburia</i> genus	TCAGACTTGCCG(C/T)ACCGC
EC1531	ND	L-S-Eco-1531-a-A-21	<i>Escherichia coli</i>	CACCGTAGTGCCCTCGTCATC

*ND, No information relating to these probes has been deposited in probeBase (<http://www.microbial-ecology.net/probebase>).

‡These probes were used together in equimolar concentrations (both at 50 ng µl⁻¹).

Formamide (20%) was included in the hybridization buffer.

doi:10.1371/journal.pone.0144457.t001

onto the surface of each well. Hybridizations were performed for 4 h in an ISO20 oven (Grant Boekel). For the washing step, slides were placed in 50 ml of wash buffer containing 20 µl of 4, 6-diamidino-2-phenylindole di-hydrochloride (DAPI; 50 ng/µl; Sigma, UK) for 15 min. They were then washed (2–3 s) in ice-cold water and dried under a stream of compressed air. Five µl of antifade reagent (polyvinyl alcohol mounting medium with DABCOTM antifading, Sigma) was added to each well and a coverslip was applied. Slides were stored in the dark at 4°C (for a maximum of 3 days) until cells were counted under a Nikon E400 Eclipse microscope. DAPI slides were visualized with the aid of a DM 400 filter and probe slides with the aid of a DM 575 filter.

The probes were commercially synthesized and labelled at the 5' end with the fluorescent dye Cy3 (Sigma-Aldrich, Poole, Dorset, UK).

Faecal moisture content

Faecal moisture content was determined following oven drying a representative sub-sample of each faecal sample in a fan-assisted oven at 100°C for 18 h.

Data transformations and statistical analysis

Data for the enumeration of the key beneficial groups of faecal bacteria were expressed as Log₁₀ cells/g fresh faeces. Within each treatment group the results for digestive tolerance and

faecal bacteriology were analysed using paired t-tests (i.e. pre-treatment vs. treatment values) within the Minitab[®] Software (Windows V15). Significance was set at $P < 0.05$.

Results

Bowel habit and indicative digestive tolerance

A summary of the mean bowel habit and indicative digestive tolerance recorded within each treatment group (i.e. 8 volunteers/treatment group) in both the 14-day pre-treatment and treatment periods is shown in Table 2. With the exception of bowel movement recordings (number/day) and stool consistency (score 1 to 7 representing hard to entirely liquid stools), volunteer responses were ranked to give rise to scores, whereby 0 indicated no symptoms, 1 indicated very mild symptoms and 10 indicated very severe symptoms. The data show that there was no significant effect ($P > 0.05$) of treatment on the mean number of bowel movements per day (1.34, 1.53, and 1.47 no/day, across respective treatments) or on stool consistency (3.7, 3.8, and 4.1, across respective treatments), values of which were similar across all treatment groups (overall mean across all treatment groups in the pre-treatment and treatment periods respectively). Determination of stool dry weight also showed that there was no significant effect ($P > 0.05$) of treatment on this between treatment groups (23.7, 28.3, and 27.1 g/100g, respectively) and that values were similar across all treatments (overall mean across all treatments 26.3, 25.6 and 26.2 g/100g fresh stool in the pre-treatment and treatment periods respectively). There was a significant effect on flatulence with the 14 g-SCF treatment, 1.33 and 2.28 in the pre-treatment and treatment periods). However, the effect of the treatment on flatulence was small, with overall values reported as none to very mild in the 14g-SCF treatment, and furthermore was not statistically significant in the 21g-SCF treatment. There was no significant effect of treatment on bloating, abdominal pain or stomach occurrences in any treatment group.

Enumeration of faecal microbial populations

Effects of the three treatments studied on predominant groups of the human gut microbiota are summarised in Table 3 and expressed as Log_{10} cells/g fresh faeces. FISH analysis showed

Table 2. Summary of the mean stool frequency and consistency, and indicative digestive tolerance recorded daily by eight volunteers per treatment group for each of 14 days in the pre-treatment (baseline) and treatment periods. Values in parenthesis are standard error (SE) values.

Parameter	8 g-SCF (6g/fibre/delivered/d)		14 g-SCF (12g/fibre/delivered/d)		21 g-SCF (18g/fibre/delivered/d)	
	Pre-treatment	Treatment	Pre-treatment	Treatment	Pre-treatment	Treatment
Bowel movements (no./day)	1.34 (0.13)	1.34 (0.14)	1.43 (0.16)	1.53 (0.15)	1.35 (0.21)	1.47 (0.16)
Stool consistency ¹	3.6 (0.20)	3.7 (0.20)	3.7 (0.26)	3.8 (0.15)	3.8 (0.22)	4.1 (0.22)
Bloating ²	0.51 (0.27)	0.89 (0.60)	0.12 (0.11)	0.42 (0.17)	1.15 (0.52)	0.93 (0.31)
Abdominal pain ²	0.77 (0.46)	0.61 (0.46)	0.05 (0.04)	0.23 (0.16)	0.98 (0.47)	0.71 (0.28)
Flatulence ²	1.54 (0.41)	1.84 (0.66)	1.33 (0.36)	2.28 (0.38)**	1.46 (0.31)	2.50 (0.78)
Stomach noises ²	0.73 (0.25)	0.92 (0.45)	0.40 (0.15)	0.96 (0.37)	0.49 (0.20)	1.30 (0.67)
Stool dry weight (g/100 g) ³	26.3 (2.71)	23.7 (1.98)	25.6 (3.37)	28.3 (2.08)	26.2 (2.00)	27.1 (1.63)

Promitor™ Soluble Corn Fibre (SCF) 8, 14 and 21 g-SCF, treatments designed to provide 6, 12 and 18 g fibre/delivered/day from SCF.

¹, estimated using the Bristol stool chart (scale with seven stool types; Type 1 –hard to Type 7 –entirely liquid).

², overall scale from 0 to 10 (0—no symptoms; 1 –very mild symptoms to 10 –very severe symptoms).

³,stool dry weight determined following oven drying at 100°C for 18 h.

*, $P < 0.05$.

** , $P < 0.01$ (vs pre-treatment).

doi:10.1371/journal.pone.0144457.t002

that there was a significant increase ($P < 0.05$) in concentration of health-promoting genus *Bifidobacterium* (Bif164) following consumption of the 6 g fibre/delivered/day when compared to the pre-treatment period. However, there was no significant increase ($P > 0.05$) in numbers of bifidobacteria for any of the other treatment groups, although the 18 g/ fibre/delivered/day indicated a trend ($P = 0.083$) for an increase (mean 9.41 and 9.65 Log_{10} cells/g fresh faeces in the pre-treatment and treatment periods respectively). There was no significant effect ($P > 0.05$) on lactobacilli although there was a trend ($P = 0.09$) for values to increase following 14-d consumption of the 6 g/ fibre/delivered/day (mean 8.07 and 8.37 Log_{10} cells/g fresh faeces in the pre-treatment and treatment periods respectively). A significant decrease ($P < 0.05$) in numbers of *C. histolyticum* /*perfringens* group following consumption of the 12 g/ fibre/delivered/day was observed (mean 7.72 and 7.19 Log_{10} cells/g fresh faeces in the pre-treatment and treatment periods respectively). *Eubacterium rectale*-*Clostridium coccoides* group enumerated by the Erec482 probe did not significantly influence ($P > 0.05$) their concentrations. A similar behaviour was demonstrated for *Bacteroides*-*Prevotella* group (detected by Bac303), Rrec584 probe enumerating the butyrate-producing *E. rectale*-*Roseburia* group, also a component of cluster XIVa, the *Atopobium* group (detected by Ato291) comprises bacteria of the *Coriobacteriaceae* family, belonging to *Coriobacterium*, *Atopobium* and *Collinsella* genera and *E. coli* (detected by Ecoli1351), whose concentrations were stably maintained during the study.

Discussion

Bowel habits and indicative digestive tolerance

The results of this ‘pilot’ study has demonstrated that SCF was well tolerated at daily dose levels of 6, 12 and 18 g/fibre/delivered/d following 14-day dietary interventions in healthy adults. Overall volunteer responses for stomach/intestinal bloating, abdominal pain, flatulence and stomach noises were recorded as none to mild (i.e. overall score < 4). These results are in agreement with previous studies [32, 33, 41, 42, 43, 44] on the digestive tolerance of prebiotics, including those determined in a single amount up to 65 g fibre/d [45]. The authors also

Table 3. Enumeration of faecal microbial populations in fresh stool samples collected from eight volunteers per treatment group both before (pre-treatment/baseline) and following 14-day dietary intervention (treatment) in a double-blind, randomised, parallel study. Stool bacterial numbers as determined by fluorescence *in situ* hybridisation are expressed as mean Log_{10} cells/g fresh faeces (values in parenthesis are standard error (SE) values).

Bacterial group	Probe	8 g-SCF		14 g-SCF		21 g-SCF	
		Pre-treatment	Treatment†	Pre-treatment	Treatment	Pre-treatment	Treatment
Total bacteria	Eub338 I-II-III	11.0 (0.07)	11.0 (0.09)	11.1 (0.08)	11.1 (0.12)	11.1 (0.10)	11.2 (0.06)
<i>Bacteroides/Prevotella</i> gp.	Bac303	10.1 (0.05)	10.2 (0.07)	10.4 (0.11)	10.2 (0.06)	10.3 (0.07)	10.4 (0.08)
Lactobacilli/enterococci	Lab158	8.07 (0.20)	8.37 (0.17)	7.87 (0.23)	7.71 (0.33)	8.08 (0.12)	8.07 (0.20)
<i>Bifidobacterium</i> spp.	Bif164	9.25 (0.27)	9.73 (0.22)*	9.26 (0.26)	9.35 (0.43)	9.41 (0.11)	9.65 (0.09)
<i>Coriobacteriaceae</i>	Ato291	9.43 (0.10)	9.37 (0.08)	9.47 (0.16)	9.39 (0.17)	9.46 (0.12)	9.52 (0.08)
<i>C. histolyticum</i> / <i>perfringens</i> gp.	Chis150	7.34 (0.15)	7.22 (0.14)	7.72 (0.16)	7.19 (0.22) *	7.51 (0.33)	7.59 (0.31)
<i>Clostridium coccoides</i> - <i>Eubacterium rectale</i> gp.	Erec482	10.3 (0.09)	10.3 (0.08)	10.4 (0.06)	10.4 (0.07)	10.4 (0.07)	10.3 (0.08)
<i>Eubacterium rectale</i> - <i>Roseburia</i> sub-gp.	Rrec584	9.87 (0.19)	9.97 (0.11)	10.0 (0.08)	9.95 (0.12)	10.0 (0.08)	9.94 (0.10)
<i>Escherichia coli</i>	Ecoli1351	8.05 (0.28)	8.00 (0.23)	7.68 (0.16)	7.68 (0.20)	10.0 (0.08)	9.94 (0.10)

†Treatments with 8, 14 and 21 g-SCF designed to provide 6, 12 and 18 g fibre/delivered/day from Promitor™ Soluble Corn Fibre (SCF).

*, $P < 0.05$ (vs pre-treatment).

doi:10.1371/journal.pone.0144457.t003

concluded that SCF was well tolerated at daily amounts as high as 65g [45], whereas some candidate prebiotic fibres have been shown to induce a mild laxation effect, increasing stool frequency and weight when consumed at high doses (>20 g/day)

In addition to the high digestive tolerance afforded by SCF, the present study showed that treatments were well tolerated as seen by no significant adverse effect on bowel habits, as determined by stool frequency and consistency, with similar values recorded both before and following 14-days of intervention. The values of 1.41 and 1.46 bowel movements per day recorded across all treatments in the pre-treatment and treatment periods are within the range previously reported [12–13]. For adults across Western countries, these authors state that the modal stool frequency is 1 per day; for the population as a whole the range is 3 stools per day to 3 stools per week [12–13]. Stool consistency was unchanged by dietary intervention and it is interesting to note that there was close agreement in the findings of the subjective assessment of stool consistency using the Bristol stool chart (recorded daily across 14 days during the pre-treatment and treatment periods) and objective assessment of stool moisture content (i.e. oven drying at 100°C for 18 h). Determination of stool consistency is an important attribute in the overall assessment of digestive tolerance and is related to stool moisture content (normal range, 70–80 g moisture/100 g fresh stool [13]). Therefore, the present study validates the approach of recording a subjective assessment of stool consistency in bowel habit diaries for use in dietary intervention studies since it is impracticable to routinely oven dry stool samples.

Assessment of prebiotic potential

Prebiotics are defined as ‘non digestible food ingredients that are selectively metabolised by colonic bacteria that have the capacity to improve health’ [16,17]. Prebiotics, in part due to their function as a special type of soluble fibre, can contribute to the health of the general population; and a number of challenges must be addressed in order to fully realize prebiotic benefits, including the need for greater awareness of the accumulated evidence on prebiotics among policy makers [22]. According to Tuohy et al. [42], the most frequently tested prebiotics are the non-digestible oligosaccharides, such as the fructans and galactans, which confer selective increases in bifidobacteria following short feeding periods.

While many non-digestible ‘by-pass’ carbohydrate materials/products may be potential candidate prebiotics, this classification cannot be extended to all non-digestible carbohydrates and dietary fibres [16]. Key to the establishment of novel prebiotic products is assessment of an optimum dose level required to elicit changes away from a gut flora dominated by potentially harmful bacteria towards a more benign, or beneficial, composition, since there is currently no recommended daily intake of prebiotics [16]. The current study examined the effect of 3 doses of SCF (6g, 12g and 18 g fibre/delivered/day from SCF) on the human gut microbiota following a 14-d intervention. In the present study, fluorescence *in situ* hybridization (FISH) method instead of 16S sequencing was used to better focus on some key taxa of interest and to obtain numerical counts of bacterial groups that are considered relevant based on previous work. Furthermore, 16S sequencing approaches might lead to bias and has previously led to under-representation of bifidobacteria. Numbers of bifidobacteria were shown to be significantly higher for the 8 g-SCF treatment. The numerical increase of 0.5 Log₁₀ bifidobacteria/g fresh stool is consistent with numerical changes of 0.5–1.0 Log₁₀ observed for inulin-type fructans [33] which are thought to exert their prebiotic effects at doses higher than 5 g/d—usually 8 g/d. According to Kolida and Gibson [46] such a change in numbers of bifidobacteria ‘constitutes a major shift in the gut microbiota towards a “healthier” composition’. This prebiotic effect concurs with that demonstrated *in vitro*, whereby relative numbers of bifidobacteria were shown to increase [46]. In addition, results of *in vivo* and *in vitro* studies show that SCF promotes the production

of SCFA including butyrate, which are positively associated with colonic health. However, increasing levels of SCFA was generally not associated with further significant changes in numbers of bifidobacteria [47].

This is concordant with other in vivo intervention studies with fructans which often reported increases between 0.5–1.0 Log₁₀ bifidobacteria counts [46]. However, the magnitude of increase seems to depend on the baseline values and higher increases are often observed in volunteers with low initial bifidobacteria numbers. While the bifidogenic potential is well-proven the effect of consumption of SCF on other bacteria groups is less well established. A decrease in clostridia was mainly reported at the dose of 12 g/fibre/delivered/day compared to the pre-treatment. However, it cannot be determined whether the decrease in clostridia after SCF intake should be seen as detrimental.

The present study has demonstrated that the optimum prebiotic dose-response was 6 g fibre/delivered/day from SCF. A low dose of SCF was shown to stimulate bifidobacteria in the gut microbiota. However, such an increase in numbers of bifidobacteria has been demonstrated over a wide range of doses in previous prebiotic studies [47–48]. The range in dose levels has been estimated to be 3–20 g/d [33] and 4–40 g/d [34] for inulin, although for individual prebiotics, differences in optimal dose level are likely to vary according to several factors including, degree of polymerisation and molecular weight [42]. An increase in the dose of SCF greater than 6 g fibre/delivered/day was generally not associated with further stimulation of the bacterial groups included for study here. This absence of a quantifiable or defined effect with increasing prebiotic inclusion is in general agreement with previously published studies investigating prebiotic dose-response effects [49–51]. In a placebo-controlled, cross-over study (n = 30), Kolida et al. [46] showed that there was no dose-response relationship following ingestion of 8 g inulin/d. Comparing daily doses of 5, 10, 20 and 40 g gum Arabic/d [52] the authors showed that the optimum daily dose was approximately 10 g/d. The explanation for the general absence of dose-response relationships is not widely understood. One such explanation proposed by Calame et al. [52] is at high doses other bacterial strains have easier access to the substrate and subsequently, less becomes available for the ones determined within their study. This is a possibility requiring further research.

In the current study numbers of faecal lactobacilli/enterococci were unchanged compared to pre-treatment values although a mean numerical increase of 0.3 Log₁₀ cells/g fresh stool was recorded for the 6 g/fibre/delivered/day as compared to the pre-treatment. Since no significant effect of dietary intervention was observed on numbers of total bacteria results of the present study suggest that consumption of 6 g/ fibre/delivered/day from SCF is selectively fermented by *Bifidobacterium* spp. and that their relative contribution to the overall gut microbiota is thereby increased.

In conclusion, SCF is a fibre source, which was well tolerated in doses as high as 18 g/ fibre/delivered/day in this study. A possible prebiotic effect was demonstrated by its stimulation of numbers of bifidobacteria in the overall gut microbiota following consumption of 6 g fibre/delivered/day. Further work is required to demonstrate the effect of SCF on other bacterial populations and in a placebo-controlled, cross over study—to assess specificity and therefore to support the potential application of this novel fiber as prebiotic in human nutrition.

Acknowledgments

The present study was funded by Tate & Lyle Ingredients Americas LLC. The research and all publications arising out of or referable to it are considered proprietary data to which Tate & Lyle Ingredients Americas LLC claims exclusive right of reference in accordance with Regulation (EC) no. 1924/2006 of the European Parliament and of the Council on Nutrition and Health Claims Made on Foods.

Author Contributions

Conceived and designed the experiments: AC ED GRG. Performed the experiments: ED AMM. Analyzed the data: AC ED. Contributed reagents/materials/analysis tools: AC GRG. Wrote the paper: AC GRG.

References

1. Huttenhower C (2012) Human Microbiome Project Consortium. Structure, function and diversity of the healthy human microbiome. *Nat* 486(7402):207–214
2. Ding T, Schloss PD (2014) Dynamics and associations of microbial community types across the human body. *Nat* 509(7500):357–360
3. Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, Sargent M, et al. (2005) Diversity of the human intestinal microbial flora. *Sci* 308: 1635–1638
4. Marteau P, Pochart P, Doré J, Béra-Maillet C, Bernalier A, Corthier G (2001). Comparative study of bacterial groups within the human cecal and fecal microbiota. *AEM* 67: 4939–4942
5. Druart C, Alligier M, Salazar N, Neyrinck AM, Delzenne NM (2014) Modulation of the gut microbiota by nutrients with prebiotic and probiotic properties. *Adv Nutr* 15; 5(5):624S–33S. doi: [10.3945/an.114.005835](https://doi.org/10.3945/an.114.005835) Print 2014 Sep PMID: [25225347](https://pubmed.ncbi.nlm.nih.gov/25225347/)
6. Kaiko GE, Stappenbeck TS (2014) Host-microbe interactions shaping the gastrointestinal environment. *Trends Immunol Sep* 11. pii: S1471-4906(14)00133-1. doi: [10.1016/j.it.2014.08.002](https://doi.org/10.1016/j.it.2014.08.002)
7. Turrioni F, Duranti S, Bottacini F, Guglielmetti S, Van Sinderen D, Ventura M (2014) *Bifidobacterium bifidum* as an example of a specialized human gut commensal. *Front Microbiol* 21; 5:437. doi: [10.3389/fmicb.2014.00437](https://doi.org/10.3389/fmicb.2014.00437) eCollection 2014. Review. Free PMC PMID: [25191315](https://pubmed.ncbi.nlm.nih.gov/25191315/)
8. Lin CS, Chang CJ, Lu CC, Martel J, Ojcius DM, Ko YF, et al. (2014) Impact of the gut microbiota, prebiotics, and probiotics on human health and disease. *Biomed J Sep* 2. doi: [10.4103/2319-4170.138314](https://doi.org/10.4103/2319-4170.138314)
9. Candela M, Biagi E, Turrioni S, Maccaferri S, Figini P, Brigidi P (2013) Dynamic efficiency of the human intestinal microbiota. *Crit Rev Microbiol. Sep* 9:1–7
10. Maukonen J, Saarela M (2014) Human gut microbiota: does diet matter? *Proc Nutr Soc* 26: 1–14
11. Konstantinidis K, Tiedje JM (2007) Prokaryotic taxonomy and phylogeny in the genomic era: advancements and challenges ahead. *Curr Opin Microbiol* 5: 504–509
12. Cummings JH, Antoine JM, Azpiroz F, Bourdet-Sicard R, Brandtzaeg P, Calder PC, et al. (2004) Gut Health and Immunity. *Eur J Nutr* 43 Suppl 2, II118–II173
13. Cummings JH, Macfarlane GT (1991) The control and consequences of bacterial fermentation in the human colon. *J Appl Bacteriol* 70(6): 443–59
14. Macfarlane GT, Gibson GR, Cummings JH (1992) Comparison of fermentation reactions in different regions of the human colon. *J Appl Bacteriol* 72(1): 57–64 PMID: [1541601](https://pubmed.ncbi.nlm.nih.gov/1541601/)
15. Slavin JL (1987) Dietary fiber: Classification, chemical analyses, and food sources. *J Am Diet Assoc* 87: 1164–1171 PMID: [3040839](https://pubmed.ncbi.nlm.nih.gov/3040839/)
16. Gibson GR, Roberfroid MB (1995) Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J Nutr* 125:1401–1412 PMID: [7782892](https://pubmed.ncbi.nlm.nih.gov/7782892/)
17. Roberfroid MB (2008) Prebiotics: concept, definition, criteria, methodologies and products. In *Handbook of Prebiotics* edited by Gibson GR and Roberfroid MB, CRC Press, Abingdon, Oxon, UK, p. 39–68
18. Hill C, Guarner F, Reid G, Gibson GR, Merenstein DJ, Pot B, et al. (2014) Expert consensus document. The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol. Aug*; 11 (8):506–14. doi: [10.1038/nrgastro.2014.66](https://doi.org/10.1038/nrgastro.2014.66) Epub 2014 Jun 10 PMID: [24912386](https://pubmed.ncbi.nlm.nih.gov/24912386/)
19. Sanders ME, Lenoir-Wijnkoop I, Salminen S, Merenstein DJ, Gibson GR, Petschow BW, et al. (2014) Probiotics and prebiotics: prospects for public health and nutritional recommendations. *Ann N Y Acad Sci. Feb*; 1309:19–29. doi: [10.1111/nyas.12377](https://doi.org/10.1111/nyas.12377) PMID: [24571254](https://pubmed.ncbi.nlm.nih.gov/24571254/)
20. Van Loveren H, Sanz Y, Salminen S (2012) Health claims in Europe: probiotics and prebiotics as case examples. *Annu Rev Food Sci Technol* 3: 247–261 doi: [10.1146/annurev-food-022811-101206](https://doi.org/10.1146/annurev-food-022811-101206) PMID: [22224554](https://pubmed.ncbi.nlm.nih.gov/22224554/)
21. Guarner F, Khan AG, Garisch J, Eliakim R, Gangl A, Thomson A, et al. (2011) World Gastroenterology Organisation Global Guidelines: probiotics and prebiotics *J Clin Gastroenterol Jul*; 46(6):468–81. doi: [10.1097/MCG.0b013e3182549092](https://doi.org/10.1097/MCG.0b013e3182549092)

22. Walsh CJ, Guinane CM, O'Toole PW, Cotter PD (2014) Beneficial modulation of the gut microbiota. *FEBS Lett.* 2014 Nov 17; 588(22):4120–4130 doi: [10.1016/j.febslet.2014.03.035](https://doi.org/10.1016/j.febslet.2014.03.035) PMID: [24681100](https://pubmed.ncbi.nlm.nih.gov/24681100/)
23. Hold GH, Smith M, Grange C, Watt RE, El-Omar EM, Mukhopadhyay I (2014) Role of the gut microbiota in inflammatory bowel disease pathogenesis: What have we learnt in the past 10 years? *World J Gastroenterol* 7; 20(5): 1192–1210 doi: [10.3748/wjg.v20.i5.1192](https://doi.org/10.3748/wjg.v20.i5.1192) PMID: [24574795](https://pubmed.ncbi.nlm.nih.gov/24574795/)
24. Golowczyc MA, Mobili P, Garrote GL, Abraham AG, De Antoni GL (2007) Protective action of *Lactobacillus kefir* carrying S-layer protein against *Salmonella enterica* serovar Enteritidis. *Int J Food Microbiol* 118(3): 264–73 PMID: [17719671](https://pubmed.ncbi.nlm.nih.gov/17719671/)
25. Mann GV, Spoerry A (1974) Studies of a surfactant and cholesteremia in the Maasai. *Am J Clin Nutr* 27: 464–469 PMID: [4596028](https://pubmed.ncbi.nlm.nih.gov/4596028/)
26. Goel AK, Dilbaghi N, Kamboj DV, Singh L (2006) Probiotics: Microbial therapy for health modulation *Def Sci J* 56(4): 513–529
27. Boge T, Rémygny M, Vaudaine S, Tanguy J, Bourdet-Sicard R, van der Werf SA (2009) Probiotic fermented dairy drink improves antibody response to influenza vaccination in the elderly in two randomised controlled trials. *Vaccine*. 27(41): 5677–84 doi: [10.1016/j.vaccine.2009.06.094](https://doi.org/10.1016/j.vaccine.2009.06.094) PMID: [19615959](https://pubmed.ncbi.nlm.nih.gov/19615959/)
28. Strozzi GP, Mogna L (2008) Quantification of folic acid in human feces after administration of *Bifidobacterium* probiotic strains. *J Clin Gastroenterol* 42(3): S179–84
29. Sangwan V, Tomar SK, Singh RRB, Singh AK, Ali B (2011) Galactooligosaccharides: novel components of designer foods. *J Food Sci* 76: R103–R111 doi: [10.1111/j.1750-3841.2011.02131.x](https://doi.org/10.1111/j.1750-3841.2011.02131.x) PMID: [22417365](https://pubmed.ncbi.nlm.nih.gov/22417365/)
30. Torres DPM, Goncalves MDF, Teixeira JA, Rodrigues LR (2010) Galacto-oligosaccharides: production, properties, applications, and significance as prebiotics. *Comprehensive Reviews in Food Science and Food Safety* 9: 438–454
31. Maathuis A, Hoffman A, Evans A, Sanders L, Venema K (2009) The effect of maize based fibers on the activity and composition of the microbiota determined in a dynamic in vitro model of the human large intestine. *J A Coll Nutr* 28(6): 657–66
32. Vester Boler BM, Rossoni Sero MC, Bauer LL, Staeger MA, Boileau TW, Swanson KS, et al. (2011) Digestive physiological outcomes related to polydextrose and soluble maize fibre consumption by healthy adult men. *Br J Nutr* 1–8
33. Konings E, Schoffelen PF, Stegen J, Blaak EE (2013) Effect of polydextrose and soluble maize fibre on energy metabolism, metabolic profile and appetite control in overweight men and women. *Br J Nutr* 1–11
34. Stewart ML, Nikhanj SD, Timm DA, Thomas W, Slavin JL (2010) Evaluation of the effect of four fibers on laxation, gastrointestinal tolerance and serum markers in healthy humans. *Ann Nutr Metab* 56: 91–98 doi: [10.1159/000275962](https://doi.org/10.1159/000275962) PMID: [20090313](https://pubmed.ncbi.nlm.nih.gov/20090313/)
35. Bassaganya-Riera J, DiGuardo M, Viladomiu M, de Horna A, Sanchez S, Einerhand AWC, et al. (2011) Soluble fibers and resistant starch ameliorate disease activity in an experimental model of inflammatory bowel disease. *J Nutr* 141(7):1318–25 doi: [10.3945/jn.111.139022](https://doi.org/10.3945/jn.111.139022) PMID: [21562241](https://pubmed.ncbi.nlm.nih.gov/21562241/)
36. Whisner CM, Martin BR, Nakatsu CH, McCabe GP, McCabe LD, Peacock M, et al. (2014) Soluble maize fibre affects short-term calcium absorption in adolescent boys and girls: a randomised controlled trial using dual stable isotopic tracers. *Br J Nutr* 112:446–456 doi: [10.1017/S0007114514000981](https://doi.org/10.1017/S0007114514000981) PMID: [24848974](https://pubmed.ncbi.nlm.nih.gov/24848974/)
37. Timm DA, Thomas W, Boileau TW, Williamson-Hughes PS, Slavin JL (2013) Polydextrose and soluble corn fiber increase five-day fecal wet weight in healthy men and women. *J Nutr* 143:473–478 doi: [10.3945/jn.112.170118](https://doi.org/10.3945/jn.112.170118) PMID: [23427334](https://pubmed.ncbi.nlm.nih.gov/23427334/)
38. Boler BM, Sero MC, Bauer LL, Staeger MA, Boileau TW, Swanson KS, et al. (2011) Digestive physiological outcomes related to polydextrose and soluble maize fibre consumption by healthy adult men. *Br J Nutr* 106:1864–71 doi: [10.1017/S0007114511002388](https://doi.org/10.1017/S0007114511002388) PMID: [21736814](https://pubmed.ncbi.nlm.nih.gov/21736814/)
39. Lewis SJ, Heaton KW (1997) Stool form scale as a useful guide to intestinal transit time. *J Scand J Gastroenterol Sep*; 32(9):920–4 PMID: [9299672](https://pubmed.ncbi.nlm.nih.gov/9299672/)
40. Costabile A, Kolida S, Klinder A, Gietl E, Bäuerlein M, Frohberg C, et al. (2010) A double-blind, placebo-controlled, cross-over study to establish the bifidogenic effect of a very-long-chain inulin extracted from globe artichoke (*Cynara scolymus*) in healthy human subjects. *Br J Nutr* 104(7):1007–17 doi: [10.1017/S0007114510001571](https://doi.org/10.1017/S0007114510001571) PMID: [20591206](https://pubmed.ncbi.nlm.nih.gov/20591206/)
41. Adam-Perrot A, Gutton L, Sanders L, Bouvier S, Combe C, Van Den Abbeele R, et al. (2009) Resistant starch and starch derived oligosaccharides as prebiotics. In *Prebiotics and Probiotics Science and Technology* 9: 259–293

42. Tuohy KM, Rouzaud GCM, Bruck WM, Gibson GR (2005) Modulation of the human gut microflora towards improved health using prebiotics—assessment of efficacy. *Curr Pharma Des* 11:75–90
43. Housez B, Cazaubiel M, Vergara C, Bard JM, Adam A, Einerhand A, et al. (2012) Evaluation of digestive tolerance of a soluble corn fibre. *J Hum Nutr Diet Oct*; 25(5):488–96. doi: [10.1111/j.1365-277X.2012.01252.x](https://doi.org/10.1111/j.1365-277X.2012.01252.x) Epub 2012 Jun 6 PMID: [22672058](https://pubmed.ncbi.nlm.nih.gov/22672058/)
44. Timm DA, Thomas W, Boileau TW, Williamson-Hughes PS, Slavin JL (2013) Polydextrose and soluble corn fiber increase five-day fecal wet weight in healthy men and women. *J Nutr.* 2013 Apr; 143(4):473–8. doi: [10.3945/jn.112.170118](https://doi.org/10.3945/jn.112.170118) Epub 2013 Feb 20 PMID: [23427334](https://pubmed.ncbi.nlm.nih.gov/23427334/)
45. Housez B, Cazaubiel M, Vergara C, Bard JM, Adam A, Einerhand A, et al. (2012) Evaluation of digestive tolerance of a soluble corn fibre. *J Hum Nutr Diet* 25(5):488–96. doi: [10.1111/j.1365-277X.2012.01252.x](https://doi.org/10.1111/j.1365-277X.2012.01252.x) Epub 2012 Jun 6 PMID: [22672058](https://pubmed.ncbi.nlm.nih.gov/22672058/)
46. Kolida S, Gibson GR (2007) Prebiotic capacity of inulin-type fructans. *J Nutr* 137: 2503S–2506S PMID: [17951493](https://pubmed.ncbi.nlm.nih.gov/17951493/)
47. Maathuis AI, Hoffman A, Evans A, Sanders L, Venema K (2009) The effect of the undigested fraction of maize products on the activity and composition of the microbiota determined in a dynamic in vitro model of the human proximal large intestine. *J Am Coll Nutr* 28(6): 657–66 PMID: [20516265](https://pubmed.ncbi.nlm.nih.gov/20516265/)
48. Whisner CM, Martin BR, Nakatsu CH, McCabe GP, McCabe LD, Peacock M, et al. (2014) Soluble maize fibre affects short-term calcium absorption in adolescent boys and girls: a randomised controlled trial using dual stable isotopic tracers. *Br J Nutr Aug*; 112(3):446–56. doi: [10.1017/S0007114514000981](https://doi.org/10.1017/S0007114514000981) Epub 2014 May 22 PMID: [24848974](https://pubmed.ncbi.nlm.nih.gov/24848974/)
49. Hooda S, Boler BM, Seroo MC, Brulc JM, Staeger MA, Boileau TW, et al. (2012) 454 pyrosequencing reveals a shift in fecal microbiota of healthy adult men consuming polydextrose or soluble corn fiber. *J Nutr Jul*; 142(7):1259–65. doi: [10.3945/jn.112.158766](https://doi.org/10.3945/jn.112.158766) Epub 2012 May 30 PMID: [22649263](https://pubmed.ncbi.nlm.nih.gov/22649263/)
50. Bouhnik Y, Raskine L, Simoneau G, Vicaut E, Neut C, Flourié B, et al. (2004) The capacity of nondigestible carbohydrates to stimulate fecal bifidobacteria in health humans: a double-blind, randomised, placebo-controlled, parallel-group, dose-response relation study. *AM J Clin Nutr* 80: 1658–1664 PMID: [15585783](https://pubmed.ncbi.nlm.nih.gov/15585783/)
51. Kolida S, Meyer D, Gibson GR (2007) A double-blind placebo-controlled study to establish the bifidogenic dose of inulin in health humans. *Eur J Clin Nutr* 61:1189–1195 PMID: [17268410](https://pubmed.ncbi.nlm.nih.gov/17268410/)
52. Calame W, Weseler AR, Viebke C, Flynn C, Siemensma AD (2008) Gum arabic establishes prebiotic functionality in healthy human volunteers in a dose-dependent manner. *Br J Nutr* 100:1269–1275 doi: [10.1017/S0007114508981447](https://doi.org/10.1017/S0007114508981447) PMID: [18466655](https://pubmed.ncbi.nlm.nih.gov/18466655/)